



❑ In addition to the tank-mix recommendations on page 1, Prosaro can also be tank-mixed with one additional product from either of the lists of insecticides or micro-nutrients.

❑ **Do not mix Prosaro + Hussar with micro-nutrients or insecticides. Do not mix with Platform S.**

Insecticides	Micro-nutrients			
Aphox	Headland Mn 15%	Verdi-Crop 4 Yield Extra*	YaraVita Magphos K	YaraVita Photrel
Cypermethrin	Headland Respond	Verdi-Crop Zinic	YaraVita Mancozin	YaraVita Phosamco
Decis	Headland Spirit	YaraVita Coptrel 500	YaraVita ManCuflo	YaraVita Zintrac 700
Fury 10 EW	Verdi-Crop Human Xtra	YaraVita Croplift	YaraVita Mantrac DF	
Hallmark	Verdi-Crop Magnor	YaraVita Foliar Potash	YaraVita Mantrac 500	
Mavrik	Verdi-Crop Sitec	YaraVita Liquid Mantrac 15%	YaraVita Sulpur F3000	
Sumi-Alpha	Verdi-Crop Supha N	YaraVita Magflo 300	YaraVita Stopit	

* Do not apply in less than 100L of water/ha

Additional 2/3 way tank mixes		
Prosaro + Axial + Adigor (apply Adigor last)	Prosaro + Cyflamid	Prosaro + Stronghold
Proline + Canopy	Prosaro + Flexity	Prosaro + Swipe
	Prosaro + Jenton	Prosaro + Talius

Tank-mix precautions

PGR tank mixtures

Prosaro + strobilurin may be applied, before crop GS32, in mixture with one of the following (dose/ha): -

Moddus 0.2L + chlormequat 1.25L (winter wheat)

Moddus 0.1L + chlormequat 1.25L (winter barley – light soils or malting crops)

Moddus 0.2L + chlormequat 1.25L (winter barley – Other soils and feed crops)

Before using any of the mixtures shown above, first check the manufacturer's latest recommendations for Moddus. DO NOT use Prosaro + strobilurin + Moddus + chlormequat on Spring Barley.

Adjuvants

In soft crops avoid the use of tank-mixtures with HBNs, Atlantis WG + biopower or Pacifica + biopower. In particular, tank mixtures with EC fungicide formulations as this may result in transitory adverse crop effects.

Sulphonyl Ureas (SU)

All sulphonyl urea herbicides have the potential to cause damage to sensitive crops if residues are left in the spray tank. For this reason sprayer hygiene following the use of SU + chlorothalonil + Prosaro is very important. Sprayers should always be thoroughly cleaned out with a proprietary sprayer cleaner immediately after use. This should always take place, even if further cereal crops have been treated by the sprayer between the last application SU mixtures and a subsequent broad leaved crop.

Cheetah Super, Grasp or Topik

Adjuvant oil recommended by the herbicide manufacturer may be added in mixes with Prosaro + strobilurin. Do NOT use any of these herbicides in mixes with Prosaro + strobilurin + Moddus + chlormequat

Trace Elements

These products are physically compatible only, there is no information on crop safety available

General Precautions

For any tank-mix it is important that the appropriate manufacturer's recommendations should be consulted and followed, and care taken to ensure that applications are made within the recommendations for each product.

First disperse Prosaro in the spray tank (follow label 'Mixing' instructions), then add the other product(s), unless specified otherwise. All mixes should be sprayed immediately and agitated throughout until the spray tank is empty. The user should ensure that all products to be tank-mixed with Prosaro are approved for use on the crops which are to be sprayed. Always check individual product labels for any possible restrictions before making a tank-mix recommendation.

Contact Bayer CropScience Limited for full details.

* Recent Changes	
14/02/2007	Canopy (PGR) tank-mix included
19/01/2007	Cyflamid + ctl and Talius + ctl included in matrix
12/12/2013	Atlantis WG tank-mix removed
13/08/2014	Format update
05/05/2020	Addition of Arizona

